

Internal number:  
 \_\_\_\_\_ / 20\_\_\_\_ / \_\_\_\_\_

**DRUG ORDER FORM**

TO BE COMPLETED BY ORDERING PHYSICIAN	<b>TREATING PHYSICIAN (stamp):</b> _____ / ____ / ____ <b>Name:</b> _____ <b>Address:</b> _____ <b>GLN Nr.:</b> _____ <b>Tel.-Nr. for inquiries:</b> _____ <div style="text-align: right;"><b>Signature</b></div>	
	The shipment and billing of medications not registered in Switzerland is <u>NOT</u> made to the patient, but <u>ALWAYS</u> to the ordering institution (hospital) or physician (office).	
	<b>DELIVERY ADDRESS:</b> _____ _____ _____	<b>BILLING ADDRESS (if deviating from delivery address):</b> _____ _____ _____
	<b>PATIENT:</b> Surname/Given name: _____ <input type="checkbox"/> M / <input type="checkbox"/> F Migration background: <input type="checkbox"/> No / <input type="checkbox"/> Yes ⇒ Country of origin: _____ Date of birth (DD/MM/JJJJ): ____ / ____ / _____ Body weight: _____ kg	
	<b>INDICATION / DIAGNOSIS:</b> _____ <b>INFECTION (MOST LIKELY) ACQUIRED IN:</b> _____	
	<b>ORDERED DRUG:</b> _____ _____	

**Note on Praziquantel:** Patients with migration background from countries endemic for neurocysticercosis should be screened by serology for the presence of asymptomatic infection before receiving praziquantel (-> risk to induce seizures in patients with asymptomatic infection) -> to order serology for "T. solium (Zystizerkose)" -> "AK", visit our laboratory website:

<https://www.swisstph.ch/en/activities/diagnostic-centre/diagnostic-orders/>

**Note on Primaquine:** Primaquine can induce severe haemolysis in patients with G6PD-deficiency! -> Assessing G6PD-status before administering primaquine is mandatory (testing for G6PD is not available from the diagnostic laboratory of the Swiss TPH).

 Send completed form via fax (061-284-8183) or via email (tropmed@swisstph.ch). Incompletely filled forms cannot be processed!

TO BE COMPLETED BY TROPICAL MEDICINE PHYSICIAN	<b>RECOMMENDED TREATMENT REGIMEN / DOSAGE:</b> _____ / ____ / ____
	_____ <b>Signature</b>